

# PRV

PATENT- OCH REGISTRERINGSVERKET

Patentavdelningen

**Intyg  
Certificate**

REC'D 03 NOV 2004

WIPO

PCT

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

*This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.*



(71) Sökande Biovitrum AB, Stockholm SE  
Applicant (s)

(21) Patentansökningsnummer 0302760-4  
Patent application number

(86) Ingivningsdatum 2003-10-20  
Date of filing

Stockholm, 2004-10-27

För Patent- och registreringsverket  
For the Patent- and Registration Office

  
Görel Gustafsson

Avgift  
Fee

**PRIORITY DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)

**PATENT- OCH  
REGISTRERINGSVERKET  
SWEDEN**

Postadress/Adress  
Box 5055  
S-102 42 STOCKHOLM

Telefon/Phone  
+46 8 782 25 00  
Vx 08-782 25 00

Telex  
17978  
PATOREG S

Telefax  
+46 8 666 02 86  
08-666 02 86

## NEW COMPOUNDS

## TECHNICAL FIELD

- 5        The present invention relates to novel compounds, to pharmaceutical compositions comprising the compounds, to processes for their preparation, as well as to the use of the compounds for the preparation of a medicament against 5-HT<sub>6</sub> receptor-related disorders.

## BACKGROUND OF THE INVENTION

10

Obesity is a condition characterized by an increase in body fat content resulting in excess body weight above accepted norms. Obesity is the most important nutritional disorder in the western world and represents a major health problem in all industrialized countries. This disorder leads to increased mortality due to increased incidences of diseases  
15 such as cardiovascular disease, digestive disease, respiratory disease, cancer and type 2 diabetes. Searching for compounds, which reduce body weight has been going on for many decades. One line of research has been activation of serotonergic systems, either by direct activation of serotonin receptor subtypes or by inhibiting serotonin reuptake. The exact receptor subtype profile required is however not known.

20

Serotonin (5-hydroxytryptamine or 5-HT), a key transmitter of the peripheral and central nervous system, modulates a wide range of physiological and pathological functions, including anxiety, sleep regulation, aggression, feeding and depression. Multiple serotonin receptor subtypes have been identified and cloned. One of these, the 5-HT<sub>6</sub> receptor, was cloned by several groups in 1993 (Ruat, M. et al. (1993) Biochem. Biophys. Res. Commun. 193: 268-276; Sebben, M. et al. (1994) NeuroReport 5: 2553-2557). This  
25 receptor is positively coupled to adenylyl cyclase and displays affinity for antidepressants such as clozapine. Recently, the effect of 5-HT<sub>6</sub> antagonist and 5-HT<sub>6</sub> antisense oligonucleotides to reduce food intake in rats has been reported (Bentley, J.C. et al. (1999) Br J Pharmacol. Suppl. 126, P66; Bentley, J.C. et al. (1997) J. Psychopharmacol. Suppl. A64, 255; Woolley M.L. et al. (2001) Neuropharmacology 41: 210-219).

30

Compounds with enhanced affinity and selectivity for the 5-HT<sub>6</sub> receptor have been identified, e.g. in WO 00/34242 and by Isaac, M. et al. (2000) *6-Bicyclopiperazinyl-1-arylsulphonylindoles and 6-Bicyclopiperidinyl-1-arylsulphonylindoles derivatives as novel,*

*potent and selective 5-HT<sub>6</sub> receptor antagonists*. Bioorganic & Medicinal Chemistry Letters 10: 1719-1721 (2000), Bioorganic & Medicinal Chemistry Letters 13: 3355-3359 (2003), Expert Opinion Therapeutic Patents 12(4) 513-527 (2002).

5 It has surprisingly been found that the compounds according to the present invention show affinity for the 5-HT<sub>6</sub> receptor as antagonists at nanomolar range. Compounds according to the present invention and their pharmaceutically acceptable salts have 5-HT<sub>6</sub> receptor antagonist, agonist and partial agonist activity and are believed to be of potential use in the treatment or prophylaxis of obesity and type 2 diabetes, to achieve reduction of body weight and of body weight gain, as well as in the treatment or prophylaxis of  
10 disorders of the central nervous system such as anxiety, depression, panic attacks, memory disorders, cognitive disorders, epilepsy, sleep disorders, migraine, anorexia, bulimia, binge eating disorders, obsessive compulsive disorders, psychoses, Alzheimer's disease, Parkinson's disease, Huntington's chorea and/or schizophrenia, panic attacks, Attention Deficit Hyperactive Disorder (ADHD), withdrawal from drug abuse, neurodegenerative  
15 diseases characterized by impaired neuronal growth, and pain. The reduction of body weight and of body weight gain (e.g. treating body-weight disorders) is achieved *inter alia* by reduction of food intake. As used herein, the term "body weight disorders" refers to the disorders caused by an imbalance between energy intake and energy expenditure, resulting in abnormal (e.g., excessive) body weight. Such body weight disorders include obesity.

## 20 INFORMATION DISCLOSURE

WO 99/42465 discloses sulphonamides derivatives that bind to the 5-HT<sub>6</sub> receptor and that can be used for the treatment of CNS disorders such as anxiety, depression,  
25 epilepsy, obsessive compulsive disorders, cognitive disorders, ADHD, anorexia and bulimia, schizophrenia, and drug abuse.

WO 01/32646 A1 discloses compounds that bind to the 5-HT<sub>6</sub> receptor and that are used for the treatment of CNS disorders and which *inter alia* may be used for the treatment of eating disorders.

30 WO 99/37623 A2 discloses compounds that bind to the 5-HT<sub>6</sub> receptor and that are used for the treatment of CNS disorders and which *inter alia* may be used for the treatment of eating disorders.

WO 99/42465 A3 discloses compounds that bind to the 5-HT<sub>6</sub> receptor and that are used for the treatment of CNS disorders and which inter alia may be used for the treatment of eating disorders.

EP 0 815 861 A1 discloses compounds that bind to the 5-HT<sub>6</sub> receptor and that are used for the treatment of CNS disorders.

WO 99/02502 A2 discloses compounds that bind to the 5-HT<sub>6</sub> receptor and that are used for the treatment of CNS disorders and which inter alia may be used for the treatment of eating disorders.

WO 98/27081 A1 discloses compounds that bind to the 5-HT<sub>6</sub> receptor and that are used for the treatment of CNS disorders and which inter alia may be used for the treatment of eating disorders.

EP 0701819 discloses compounds that bind to the 5-HT<sub>1D</sub> receptor and that are used for the treatment of CNS disorders and obesity.


US 6,191,141 and WO 01/12629 disclose compounds that bind to the 5-HT<sub>6</sub> receptor and that are used for the treatment of CNS disorders.

WO03/072198 disclose benzenesulphonamide derivatives for the treatment of obesity.

No publications disclose the compounds and their use according to the present invention against 5-HT<sub>6</sub> receptor-related disorders.

## DISCLOSURE OF THE INVENTION


One object of the present invention is a compound of the Formula (I)




Formula (I)

wherein:

P is selected from a substituent of Formula (II) and Formula (III);



(II)



(III)

or P may also be selected from H or C<sub>1-6</sub>-alkyl provided that R<sup>m</sup> is selected from -NH<sub>2</sub>SO<sub>2</sub>R<sup>11</sup>, -SO<sub>2</sub>NR<sup>8</sup>R<sup>11</sup> or -S(O)<sub>e</sub>R<sup>11</sup>, wherein e is 0, 1, 2 or 3;

5

v is 1 or 2;

— represents a single bond or a double bond, with the proviso that both

— represent double bonds or that both — represent single bonds;

W<sub>1</sub>, W<sub>2</sub>, W<sub>3</sub>, Z and Y are each a carbon atom; or

10 one of W<sub>1</sub>, W<sub>2</sub>, W<sub>3</sub>, Z and Y is a nitrogen atom, while the remainder being carbon atoms, provided that both — in Formula (I) represent single bonds;

U is selected from CHR<sup>4</sup>, CR<sup>4</sup> and CR<sup>4</sup>R<sup>4'</sup>, provided that when the dotted line connecting W<sub>1</sub> and U is a double bond, then U is CR<sup>4</sup>; and further provided that when the dotted line  
15 connecting W<sub>1</sub> and U is a single bond, then U is selected from CHR<sup>4</sup> and CR<sup>4</sup>R<sup>4'</sup>;

R<sup>1</sup> is selected from:

- (a) C<sub>1-6</sub>-alkyl,
- (b) C<sub>1-6</sub>-alkoxy-C<sub>1-6</sub>-alkyl,
- 20 (c) C<sub>2-6</sub>-alkenyl,
- (d) hydroxy-C<sub>1-6</sub>-alkyl,
- (e) halo-C<sub>1-6</sub>-alkyl,
- (f) aryl,
- (g) arylcarbonylmethyl,
- 25 (h) aryl-C<sub>2-6</sub>-alkenyl,
- (i) aryl-C<sub>1-6</sub>-alkyl,
- (j) C<sub>3-7</sub>-cycloalkyl,
- (k) heteroaryl,
- (l) 4-piperidinyl,

(m) N-substituted 4-piperidinyl, wherein the substituents are selected from C<sub>1-6</sub>-alkyl, aryl-C<sub>1-6</sub>-alkyl and heteroaryl-C<sub>1-6</sub>-alkyl,

(n) heteroaryl-C<sub>1-6</sub>-alkyl,


wherein any heteroaryl or aryl residue, alone or as part of another group, is optionally substituted, independently, in one or more positions with substituents having the values as defined for R<sup>m</sup> and R<sup>m'</sup>;

R<sup>m</sup> and R<sup>m'</sup> are each independently selected from:

- (a) hydrogen,
- 10 (b) halogen,
- (c) C<sub>1-6</sub>-alkyl,
- (d) hydroxy,
- (e) C<sub>1-6</sub>-alkoxy,
- (f) C<sub>2-6</sub>-alkenyl,
- 15 (g) phenyl,
- (h) phenoxy,
- (i) benzyloxy,
- (j) benzoyl,
- (k) -OCF<sub>3</sub>,
- 20 (l) -CN,
- (m) hydroxy-C<sub>1-6</sub>-alkyl,
- (n) halo-C<sub>1-6</sub>-alkyl,
- (o) -NR<sup>10</sup>R<sup>10</sup>,
- (p) -NO<sub>2</sub>,
- 25 (q) -CONR<sup>10</sup>R<sup>10</sup>,
- (r) -NHSO<sub>2</sub>R<sup>11</sup>,
- (s) -NR<sup>8</sup>COR<sup>11</sup>,
- (t) -SO<sub>2</sub>NR<sup>8</sup>R<sup>11</sup>,
- (u) -C(=O)R<sup>11</sup>,
- 30 (v) C<sub>1-6</sub>-alkoxycarbonyl,
- (w) -S(O)<sub>e</sub>R<sup>11</sup>, wherein e is 0, 1, 2 or 3,
- (x) -SCF<sub>3</sub>,
- (y) -CHF=CH<sub>2</sub>, or
- (aa) -OCF<sub>2</sub>H;


- and with the proviso that, when  $R^m$  and  $R^{m'}$  are different from hydrogen and ring A is aromatic,  $R^m$  and  $R^{m'}$  are attached to a carbon atom in Formula (I); and with the further proviso that when one of  $W_1$ ,  $W_2$  and  $W_3$  in Formula (I) is a nitrogen atom and both  $\text{-----}$  represent single bonds the said nitrogen atom is attached to  $R^m$ , wherein  $R^m$  is selected
- 5 from hydrogen or  $C_{1-4}$ -alkyl and  $v$  is 1; and with the further proviso that when  $W_1$ ,  $W_2$  and  $W_3$  in Formula (I) are each a carbon atom and both  $\text{-----}$  represent single bonds,  $R^m$  is selected from hydrogen or methyl; and with the further proviso that when  $R^m$  or  $R^{m'}$ , as substituents on ring A and B in Formula (I), are selected from phenyl, phenoxy, benzyloxy and benzoyl, the phenyl or aryl ring thereof may be optionally substituted by  $C_{1-4}$ -alkyl,
- 10 halogen,  $C_{1-4}$ -alkoxy,  $C_{1-4}$ -alkylthio, trifluoromethyl or cyano;

wherein  $R^m$  and  $R^4$  may be linked to each other to form a fused substituent of Formula (IV) provided that  $R^m$  is attached to  $W_1$ :



(IV)

15 when U is  $CR^4$  or  $CHR^4$ ,  $R^4$  is a group selected from:



wherein  $R^4$  may also be a group selected from:  $O-(C=NH)NR^8R^8$  or





wherein:

$n = 0, 1, \text{ or } 2,$

$o = 1 \text{ or } 2,$


5  $p = 1, 2, 3, \text{ or } 4,$

$r = 2 \text{ or } 3,$

$s = 1, 2 \text{ or } 3,$

$t = 0, 1, 2, \text{ or } 3;$

10 when U is  $CHR^4$ ,  $R^4$  is additionally selected from the following groups:



wherein:

$n = 0, 1 \text{ or } 2,$

$o = 1 \text{ or } 2,$

15  $p = 1, 2, 3, \text{ or } 4,$

$r = 2 \text{ or } 3,$

$s = 1, 2 \text{ or } 3;$

wherein X is selected from O,  $NR^8$  and S;

20

wherein  $R^5$  is independently a group selected from:

(a) hydrogen,

(b)  $C_{1-6}$ -alkyl,

(c) 2-cyanoethyl,

25

(d) hydroxy- $C_{2-6}$ -alkyl,

- (e) C<sub>3-6</sub>-alkenyl,  
 (f) C<sub>3-6</sub>-alkynyl,  
 (g) C<sub>3-7</sub>-cycloalkyl,  
 (h) C<sub>3-7</sub>-cycloalkyl-C<sub>1-4</sub>-alkyl,  
 5 (i) C<sub>1-6</sub>-alkoxy-C<sub>2-6</sub>-alkyl,  
 (j) aryl-C<sub>1-6</sub>-alkyl,  
 (k) heteroaryl-C<sub>1-6</sub>-alkyl,  
 (l) -C(=NH)-NR<sup>8</sup>R<sup>8</sup>,  
 (m) -C(=O)-NR<sup>8</sup>R<sup>8</sup>,  
 10 (n) -CH<sub>2</sub>-CO-NR<sup>8</sup>R<sup>8</sup>;

R<sup>6</sup> is selected from:

- (a) hydrogen,  
 (b) C<sub>1-4</sub>-alkyl,  
 15 (c) hydroxy-C<sub>1-4</sub>-alkyl,  
 (d) C<sub>1-4</sub>-alkoxy-C<sub>1-4</sub>-alkyl,  
 (e) halo-C<sub>1-4</sub>-alkyl,  
 (f) -NR<sup>8</sup>R<sup>8</sup>,  
 (g) -CO-NR<sup>8</sup>R<sup>8</sup>,  
 20 (h) hydroxy, provided that the said hydroxy group is not attached to a carbon atom adjacent to a ring nitrogen atom;

R<sup>7</sup> is selected from:

- (a) hydrogen,  
 25 (b) C<sub>1-4</sub>-alkyl,  
 (c) hydroxy-C<sub>1-4</sub>-alkyl, or  
 (d) C<sub>1-4</sub>-alkoxy-C<sub>1-4</sub>-alkyl,  
 (e) hydroxy, provided that the said hydroxy group is not attached to a carbon atom adjacent to a heterocyclic ring nitrogen atom and that the said hydroxy group is  
 30 attached to a heterocyclic ring not substituted with oxo;

R<sup>8</sup> is each independently selected from:

- (a) hydrogen, or  
 (b) C<sub>1-6</sub>-alkyl,

$R^9$  is selected from:

- (a) hydrogen;
- (b)  $C_{1-4}$ -alkyl, wherein one or two groups may be present at any carbon atom, or  
 5 when two groups are present at the same carbon atom they may together form a cyclopropane ring,
- (c) hydroxy- $C_{1-4}$ -alkyl,
- (d)  $C_{1-4}$ -alkoxy- $C_{1-4}$ -alkyl,
- (e) halo- $C_{1-4}$ -alkyl,
- 10 (f)  $-CO-NR^8R^8$ ;

$R^{10}$  is each independently selected from:

- (a) hydrogen,
- (b)  $C_{1-6}$ -alkyl
- 15 (c)  $C_{3-7}$ -cycloalkyl, or

wherein the two  $R^{10}$  groups together with the nitrogen to which they are attached form a heterocyclic ring; and when the two  $R^{10}$  groups form a piperazine ring, the nitrogen of the piperazine ring that allows the substitution may be substituted with a group selected from  $R^5$ ;

20  $R^{11}$  is selected from:

- (a)  $C_{1-6}$ -alkyl,
- (b) aryl, or
- (c) heteroaryl;


25  $R^{12}$  is selected from:

- (a) hydrogen, or
- (b) methyl;

30 when U is  $R^4R^{4'}$ ,  $R^4$  and  $R^{4'}$  are linked to each other to form a heterocyclic ring selected from pyrrolidine or piperidine, wherein the N atom may be substituted by a group selected from  $R^5$ ; and pharmaceutically acceptable salts, hydrates, solvates, geometrical isomers, tautomers, optical isomers, and prodrug forms thereof.

It is preferred in Formula (I) that:


P is selected from



(II)

- 5 each of  $W_1$ ,  $W_2$ ,  $W_3$ , Z and Y is a carbon atom provided that both  $\text{-----}$  in Formula (I) represent double bonds; or  
one of  $W_1$ ,  $W_2$ ,  $W_3$ , Z and Y is a nitrogen atom, while the remainder being carbon atoms, provided that both  $\text{-----}$  in Formula (I) represent single bonds;
  - 10 U is selected from  $\text{CHR}^4$ ,  $\text{CR}^4$  and  $\text{CR}^4\text{R}^{4'}$ , provided that when the dotted line connecting  $W_1$  and U is a double bond, then U is  $\text{CR}^4$ ; and further provided that when the dotted line connecting  $W_1$  and U is a single bond, then U is selected from  $\text{CHR}^4$  and  $\text{CR}^4\text{R}^{4'}$ ;
- $\text{R}^1$  is selected from:
- 15 (f) aryl,
  - (h) aryl- $\text{C}_{2-6}$ -alkenyl,
  - (i) aryl- $\text{C}_{1-6}$ -alkyl,
  - (j)  $\text{C}_{3-7}$ -cycloalkyl,
  - (k) heteroaryl,
  - 20 (n) heteroaryl- $\text{C}_{1-6}$ -alkyl,
- wherein any heteroaryl or aryl residue, alone or as part of another group, is optionally substituted, independently, in one or more positions with substituents having the values as defined for  $\text{R}^m$  and  $\text{R}^{m'}$ ;
- 25  $\text{R}^m$  and  $\text{R}^{m'}$  are each independently selected from:
    - (a) hydrogen,
    - (b) halogen,
    - (c)  $\text{C}_{1-6}$ -alkyl,
    - (d) hydroxy,
    - 30 (e)  $\text{C}_{1-6}$ -alkoxy,

- (f) C<sub>2-6</sub>-alkenyl,  
 (k) -OCF<sub>3</sub>,  
 (m) hydroxy-C<sub>1-6</sub>-alkyl,  
 (n) halo-C<sub>1-6</sub>-alkyl,  
 5 (o) -NR<sup>10</sup>R<sup>10</sup>,  
 (q) -CONR<sup>10</sup>R<sup>10</sup>,  
 (r) -NHSO<sub>2</sub>R<sup>11</sup>,  
 (s) -NR<sup>8</sup>COR<sup>11</sup>,  
 (t) -SO<sub>2</sub>NR<sup>8</sup>R<sup>11</sup>,  
 10 (u) -C(=O)R<sup>11</sup>,  
 (w) -S(O)<sub>e</sub>R<sup>11</sup>, wherein e is 0, 1, 2 or 3,  
 (x) -SCF<sub>3</sub>,  
 (y) -CHF=CH<sub>2</sub>, or  
 (aa) -OCF<sub>2</sub>H;
- 15 and with the proviso that, when R<sup>m</sup> and R<sup>m'</sup> are different from hydrogen and ring A is aromatic, R<sup>m</sup> and R<sup>m'</sup> are attached to a carbon atom in Formula (I); and with the further proviso that when one of W<sub>1</sub>, W<sub>2</sub> and W<sub>3</sub> in Formula (I) is a nitrogen atom and both        represent single bonds the said nitrogen atom is attached to R<sup>m</sup>, wherein R<sup>m</sup> is selected from hydrogen or C<sub>1-4</sub>-alkyl and v is 1; and with the further proviso that when W<sub>1</sub>, W<sub>2</sub> and
- 20 W<sub>3</sub> in Formula (I) are each a carbon atom and both        represent single bonds, R<sup>m</sup> is selected from hydrogen or methyl; and with the further proviso that when R<sup>m</sup> or R<sup>m'</sup>, as substituents on ring A and B in Formula (I), are selected from phenyl, phenoxy, benzyloxy and benzoyl, the phenyl or aryl ring thereof may be optionally substituted by C<sub>1-4</sub>-alkyl, halogen, C<sub>1-4</sub>-alkoxy, C<sub>1-4</sub>-alkylthio, trifluoromethyl or cyano;
- 25 when U is CR<sup>4</sup> or CHR<sup>4</sup>, R<sup>4</sup> is a group selected from:




$n = 0, 1, \text{ or } 2,$

$o = 1 \text{ or } 2,$

$p = 1, 2, 3, \text{ or } 4,$

5

when U is  $\text{CHR}^4$ ,  $\text{R}^4$  is additionally selected from the following groups:



wherein :

$n = 0, 1 \text{ or } 2,$

$o = 1 \text{ or } 2,$

5  $p = 1, 2, \text{ or } 3;$

$r = 2 \text{ or } 3,$

wherein X is selected from O,  $\text{NR}^8$  and S;

wherein  $\text{R}^5$  is independently a group selected from:

- 10 (a) hydrogen,  
 (b)  $\text{C}_{1-6}$ -alkyl,  
 (c) 2-cyanoethyl,  
 (d) hydroxy- $\text{C}_{2-4}$ -alkyl,  
 (e)  $\text{C}_{3-6}$ -alkenyl,  
 15 (h)  $\text{C}_{3-7}$ -cycloalkyl- $\text{C}_{1-4}$ -alkyl, or  
 (i)  $\text{C}_{1-4}$ -alkoxy- $\text{C}_{2-4}$ -alkyl,

$\text{R}^6$  is selected from:

- 20 (a) hydrogen,  
 (b)  $\text{C}_{1-4}$ -alkyl,  
 (f)  $-\text{NR}^8\text{R}^8,$

$\text{R}^7$  is selected from:

- 25 (a) hydrogen,  
 (b)  $\text{C}_{1-4}$ -alkyl,

(c) hydroxy-C<sub>1-2</sub>-alkyl, or

(d) C<sub>1-2</sub>-alkoxy-C<sub>1-2</sub>-alkyl,

(e) hydroxy, provided that the said hydroxy group is not attached to a carbon atom adjacent to a heterocyclic ring nitrogen atom and that the said hydroxy group is attached to a heterocyclic ring not substituted with oxo;

R<sup>8</sup> is each independently selected from:

(a) hydrogen, or

(b) C<sub>1-6</sub>-alkyl,

R<sup>9</sup> is selected from:

(a) hydrogen,

(b) C<sub>1-4</sub>-alkyl, wherein one or two groups may be present at any carbon atom, or when two groups are present at the same carbon atom they may together form a cyclopropane ring,

(c) hydroxy-C<sub>1-2</sub>-alkyl,

(d) C<sub>1-2</sub>-alkoxy-C<sub>1-2</sub>-alkyl,

(e) halo-C<sub>1-2</sub>-alkyl,

R<sup>10</sup> is each independently selected from:

(a) hydrogen,

(b) C<sub>1-6</sub>-alkyl, or


wherein the two R<sup>10</sup> groups together with the nitrogen to which they are attached form a heterocyclic ring; and when the two R<sup>10</sup> groups form a piperazine ring, the nitrogen of the piperazine ring that allows the substitution may be substituted with a group selected from R<sup>5</sup>;

R<sup>11</sup> is selected from:


(a) C<sub>1-6</sub>-alkyl

provided that P is a group selected from Formula (II) or Formula (III);





(II)



(III)

$R^{12}$  is selected from:

- (a) hydrogen, or
- (b) methyl;

when U is  $R^4R^{4'}$ ,  $R^4$  and  $R^{4'}$  are linked to each other to form a heterocyclic ring selected from pyrrolidine or piperidine, wherein the N atom may be substituted by a group  $R^5$  selected from:

- (a) hydrogen,
- (b)  $C_{1-6}$ -alkyl,
- (c) 2-cyanoethyl,
- (d) hydroxy- $C_{2-4}$ -alkyl,
- (i)  $C_{1-4}$ -alkoxy- $C_{2-4}$ -alkyl.

It is more preferred that

ring A is saturated;

$W_1$  is a nitrogen atom;

Z and Y are both carbon atoms;

$R^1$  is a group aryl or heteroaryl optionally substituted by halogen;

$R^m$  and  $R^{m'}$  are both H;

wherein  $R^4$  and  $R^{4'}$  are linked to each other to form piperidine, wherein the N atom is substituted by methyl.

It is even more preferred that

Ar is selected from phenyl, naphthyl, and thienyl, which group Ar is optionally substituted by halogen.

Preferred compounds are:

4'-Methyl-1'-(2-naphthylsulphonyl)-1',4',5',6'-tetrahydrospiro {piperidine-2,7'-pyrrolo[3,2-b]pyridine} hydrochloride,

4'-Methyl-1'-(4-bromophenylsulphonyl)-1',4',5',6'-tetrahydrospiro {piperidine-2,7'-pyrrolo[3,2-b]pyridine} hydrochloride,

5 4'-Methyl-1'-(5-bromo-2-thienylsulphonyl)-1',4',5',6'-tetrahydrospiro {piperidine-2,7'-pyrrolo[3,2-b]pyridine} hydrochloride, and

4'-Methyl-1'-(2-thienylsulphonyl)-1',4',5',6'-tetrahydrospiro {piperidine-2,7'-pyrrolo[3,2-b]pyridine} hydrochloride.

10 Another object of the present invention is a process for the preparation of a compound as mentioned above, comprising the following steps:

- a) reaction of 2-(2-ethylamino)pyrrole and 1-methylpiperazine-4-one to give 4'-methyl-1',4',5',6'-tetrahydrospiro {piperidine-2,7'-pyrrolo[3,2-b]pyridine}; and
- b) reaction of the product from step a) with an arylsulphonyl chloride in the presence of a base.

15 Another object of the present invention is a compound as mentioned above for use in therapy, especially for use in the treatment or prophylaxis of a 5-HT<sub>6</sub> receptor-related disorder, to achieve reduction of body weight and of body weight gain.

Another object of the present invention is a pharmaceutical formulation comprising a compound as mentioned above as active ingredient, in combination with a  
20 pharmaceutically acceptable diluent or carrier, especially for use in the treatment or prophylaxis of a 5-HT<sub>6</sub> receptor-related disorder, to achieve reduction of body weight and of body weight gain.

Another object of the present invention is a method for treating a human or animal subject suffering from a 5-HT<sub>6</sub> receptor-related disorder, to achieve reduction of body  
25 weight and of body weight gain. The method can include administering to a subject (e.g., a human or an animal, dog, cat, horse, cow) in need thereof an effective amount of one or more compounds of any of the formulae herein, their salts, or compositions containing the compounds or salts.

The methods delineated herein can also include the step of identifying that the subject  
30 is in need of treatment of the 5-HT<sub>6</sub> receptor-related disorder, to achieve reduction of body weight and of body weight gain. Identifying a subject in need of such treatment can be in the judgment of a subject or a health care professional and can be subjective (e.g., opinion) or objective (e.g., measurable by a test or diagnostic method).

Another object of the present invention is a method for the treatment or prophylaxis of a 5-HT<sub>6</sub> receptor-related disorder, to achieve reduction of body weight and of body weight gain, which comprises administering to a subject in need of such treatment an effective amount of a compound as mentioned above.

5 Another object of the present invention is a method for modulating 5-HT<sub>6</sub> receptor activity, which comprises administering to a subject in need of such treatment an effective amount of a compound as mentioned above.

Another object of the present invention is the use of a compound as mentioned above for the manufacture of a medicament for use in the prophylaxis or treatment of a 5-HT<sub>6</sub> 10 receptor-related disorder, to achieve reduction of body weight and of body weight gain.

The compounds as mentioned above may be agonists, partial agonists or antagonists for the 5-HT<sub>6</sub> receptor. Preferably, the compounds act as partial agonists or antagonists for the 5-HT<sub>6</sub> receptor.

Examples of 5-HT<sub>6</sub> receptor-related disorders are obesity; type II diabetes; disorders 15 of the central nervous system such as anxiety, depression, panic attacks, memory disorders, cognitive disorders, epilepsy, sleep disorders, migraine, anorexia, bulimia, binge eating disorders, obsessive compulsive disorders, psychoses, Alzheimer's disease, Parkinson's disease, Huntington's chorea, schizophrenia, attention deficit hyperactive disorder (ADHD), withdrawal from drug abuse, neurodegenerative diseases characterized by 20 impaired neuronal growth, and pain.

The compounds and compositions are useful for treating diseases, to achieve reduction of body weight and of body weight gain. The diseases include obesity; type II diabetes; disorders of the central nervous system such as anxiety, depression, panic attacks, memory disorders, cognitive disorders, epilepsy, sleep disorders, migraine, anorexia, 25 bulimia, binge eating disorders, obsessive compulsive disorders, psychoses, Alzheimer's disease, Parkinson's disease, Huntington's chorea, schizophrenia, attention deficit hyperactive disorder (ADHD), withdrawal from drug abuse, neurodegenerative diseases characterized by impaired neuronal growth, and pain. In one aspect, the invention relates to a method for treating or preventing an aforementioned disease comprising administering to 30 a subject in need of such treatment an effective amount or composition delineated herein.

Another object of the present invention is a cosmetic composition comprising a compound as mentioned above as active ingredient, in combination with a cosmetically

acceptable diluent or carrier, especially for use in the prophylaxis or treatment of a 5-HT<sub>6</sub> receptor-related disorder, to achieve reduction of body weight and of body weight gain.

### Definitions

5

The following definitions shall apply throughout the specification and the appended claims.

Unless otherwise stated or indicated, the term "C<sub>1-6</sub>-alkyl" denotes a straight or branched alkyl group having from 1 to 6 carbon atoms. Examples of said C<sub>1-6</sub>-alkyl include  
 10 methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, t-butyl and straight- and branched-chain pentyl and hexyl. For parts of the range "C<sub>1-6</sub>-alkyl" all subgroups thereof are contemplated such as C<sub>1-5</sub>-alkyl, C<sub>1-4</sub>-alkyl, C<sub>1-3</sub>-alkyl, C<sub>1-2</sub>-alkyl, C<sub>2-6</sub>-alkyl, C<sub>2-5</sub>-alkyl, C<sub>2-4</sub>-alkyl, C<sub>2-3</sub>-alkyl, C<sub>3-6</sub>-alkyl, C<sub>4-5</sub>-alkyl, etc. "Halo-C<sub>1-6</sub>-alkyl" means a C<sub>1-6</sub>-alkyl group substituted by one or more halogen atoms. Examples of said halo-C<sub>1-6</sub>-alkyl include 2-  
 15 fluoroethyl, fluoromethyl, trifluoromethyl and 2,2,2-trifluoroethyl. Likewise, "aryl-C<sub>1-6</sub>-alkyl" means a C<sub>1-6</sub>-alkyl group substituted by one or more aryl groups.

Unless otherwise stated or indicated, the term "hydroxy-C<sub>1-6</sub>-alkyl" denotes a straight or branched alkyl group that has a hydrogen atom thereof replaced with OH. Examples of said hydroxy-C<sub>1-6</sub>-alkyl include hydroxymethyl, 2-hydroxyethyl, 2-  
 20 hydroxypropyl and 2-hydroxy-2-methylpropyl.

Unless otherwise stated or indicated, the term "C<sub>1-6</sub>-alkoxy" denotes a straight or branched alkoxy group having from 1 to 6 carbon atoms. Examples of said C<sub>1-6</sub>-alkoxy include methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, iso-butoxy, sec-butoxy, t-butoxy and straight- and branched-chain pentoxy and hexoxy. For parts of the range "C<sub>1-6</sub>-  
 25 alkoxy" all subgroups thereof are contemplated such as C<sub>1-5</sub>-alkoxy, C<sub>1-4</sub>-alkoxy, C<sub>1-3</sub>-alkoxy, C<sub>1-2</sub>-alkoxy, C<sub>2-6</sub>-alkoxy, C<sub>2-5</sub>-alkoxy, C<sub>2-4</sub>-alkoxy, C<sub>2-3</sub>-alkoxy, C<sub>3-6</sub>-alkoxy, C<sub>4-5</sub>-alkoxy, etc.

Unless otherwise stated or indicated, the term C<sub>1-6</sub>-alkoxy-C<sub>1-6</sub>-alkyl denotes a  
 30 straight or branched alkoxy group having from 1 to 6 carbon atoms connected to an alkyl group having from 1 to 6 carbon atoms. Examples of said C<sub>1-6</sub>-alkoxy-C<sub>1-6</sub>-alkyl include methoxymethyl, ethoxymethyl, iso-propoxymethyl, n-butoxymethyl, t-butoxymethyl and straight- and branched-chain pentoxymethyl. For parts of the range "C<sub>1-6</sub>-alkoxy-C<sub>1-6</sub>-alkyl" all subgroups thereof are contemplated such as C<sub>1-5</sub>-alkoxy-C<sub>1-6</sub>-alkyl, C<sub>1-4</sub>-alkoxy-

C<sub>1-6</sub>-alkyl, C<sub>1-3</sub>-alkoxy-C<sub>1-6</sub>-alkyl, C<sub>1-2</sub>-alkoxy-C<sub>1-6</sub>-alkyl, C<sub>2-6</sub>-alkoxy-C<sub>1-6</sub>-alkyl, C<sub>2-5</sub>-alkoxy-C<sub>1-6</sub>-alkyl, C<sub>2-4</sub>-alkoxy-C<sub>1-6</sub>-alkyl, C<sub>2-3</sub>-alkoxy-C<sub>1-6</sub>-alkyl, C<sub>3-6</sub>-alkoxy-C<sub>1-6</sub>-alkyl, C<sub>4-5</sub>-alkoxy-C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy-C<sub>1-5</sub>-alkyl, C<sub>1-6</sub>-alkoxy-C<sub>1-4</sub>-alkyl, etc.

Unless otherwise stated or indicated, the term "C<sub>2-6</sub>-alkenyl" denotes a straight or  
 5 branched alkenyl group having from 2 to 6 carbon atoms. Examples of said C<sub>2-6</sub>-alkenyl include vinyl, allyl, 2,3-dimethylallyl, 1-butenyl, 1-pentenyl, and 1-hexenyl. For parts of the range "C<sub>2-6</sub>-alkenyl" all subgroups thereof are contemplated such as C<sub>2-5</sub>-alkenyl, C<sub>2-4</sub>-alkenyl, C<sub>2-3</sub>-alkenyl, C<sub>3-6</sub>-alkenyl, C<sub>4-5</sub>-alkenyl, etc. Likewise, "aryl-C<sub>2-6</sub>-alkenyl" means a  
 10 C<sub>2-6</sub>-alkenyl group substituted by one or more aryl groups. Examples of said aryl-C<sub>2-6</sub>-alkenyl include styryl and cinnamyl.

The term "oxo" denotes  $\text{=O}$ .

Unless otherwise stated or indicated, the term "C<sub>2-6</sub>-alkynyl" denotes a straight or  
 branched alkynyl group having from 2 to 6 carbon atoms. Examples of said C<sub>2-6</sub>-alkynyl include ethynyl, 1-propynyl, 1-butyne, and 1-hexynyl. For parts of the range "C<sub>2-6</sub>-  
 15 alkynyl" all subgroups thereof are contemplated such as C<sub>2-5</sub>-alkynyl, C<sub>2-4</sub>-alkynyl, C<sub>2-3</sub>-alkynyl, C<sub>3-6</sub>-alkynyl, C<sub>4-5</sub>-alkynyl, etc.

Unless otherwise stated or indicated, the term "C<sub>3-7</sub>-cycloalkyl" denotes a cyclic alkyl group having a ring size from 3 to 7 carbon atoms. Examples of said cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, methylcyclohexyl, and  
 20 cycloheptyl. For parts of the range "C<sub>3-7</sub>-cycloalkyl" all subgroups thereof are contemplated such as C<sub>3-6</sub>-cycloalkyl, C<sub>3-5</sub>-cycloalkyl, C<sub>3-4</sub>-cycloalkyl, C<sub>4-7</sub>-cycloalkyl, C<sub>4-6</sub>-cycloalkyl, C<sub>4-5</sub>-cycloalkyl, C<sub>5-7</sub>-cycloalkyl, C<sub>6-7</sub>-cycloalkyl, etc.

Unless otherwise stated or indicated, the term "aryl" refers to a hydrocarbon ring system having at least one aromatic ring. Examples of aryls are phenyl, pentalenyl,  
 25 indenyl, indanyl, 1,2,3,4-tetrahydronaphthyl, 1-naphthyl, 2-naphthyl, fluorenyl, anthryl, phenanthryl and pyrenyl. The aryl rings may be optionally substituted. Likewise, aryloxy refers to an aryl group bonded to an oxygen atom.





The term "heteroaryl" refers to a mono- or bicyclic aromatic ring system, only one ring need be aromatic, and the said heteroaryl moiety can be linked to the remainder of the  
 30 molecule via a carbon or nitrogen atom in any ring, and having from 5 to 10 ring atoms (mono- or bicyclic), in which one or more of the ring atoms are other than carbon, such as nitrogen, sulphur, oxygen and selenium. Examples of such heteroaryl rings include furyl, pyrrolyl, thienyl, oxazolyl, isoxazolyl, imidazolyl, thiazolyl, isothiazolyl, pyridinyl,

pyrimidinyl, pyrazinyl, chromanyl, quinazolinyl, indolyl, isoindolyl, indolinyl, isoindolinyl, indazolyl, pyrazolyl, pyridazinyl, quinolinyl, isoquinolinyl, benzofuranyl, dihydrobenzofuranyl, benzodioxolyl, benzodioxinyl, benzothienyl, benzimidazolyl, benzothiazolyl, benzothiadiazolyl, and benzotriazolyl groups. If a bicyclic heteroaryl ring is substituted, it may be substituted in any ring.

Unless otherwise stated or indicated, the term "heterocyclic" refers to a non-aromatic (i.e., partially or fully saturated) mono- or bicyclic ring system having 4 to 10 ring atoms with at least one heteroatom such as B, O, N, or S, and the remaining ring atoms are carbon. Examples of heterocyclic groups include piperidyl, tetrahydropyranyl, tetrahydrofuranyl, azepinyl, azetidiny, pyrrolidinyl, morpholinyl, imidazolinyl, thiomorpholinyl, pyranyl, dioxanyl, and piperazinyl groups. When present, the sulfur atom may be in an oxidized form (i.e., S=O or O=S=O). Examples of heterocyclic groups containing sulfur in oxidized form include octahydro[3,4b]pyrazine 6,6-dioxide and thiomorpholine 1,1-dioxide.

Unless otherwise stated or indicated, the term "halogen" shall mean fluorine, chlorine, bromine or iodine.

The term  $-S(O)_eR^{11}$ , wherein e is 0, 1, 2 or 3, has the meaning as illustrated by Formula (V) – (VIII):



The term "leaving group" refers to a group to be displaced from a molecule during a nucleophilic displacement reaction. Examples of leaving groups are iodide, bromide, chloride, methanesulphonate, hydroxy, methoxy, thiomethoxy, tosyl, or suitable protonated forms thereof (e.g.,  $H_2O$ ,  $MeOH$ ), especially bromide and methanesulphonate.

"Optional" or "optionally" means that the subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not.

"Pharmaceutically acceptable" means being useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable and includes being useful for veterinary use as well as human pharmaceutical use.

"Treatment" as used herein includes prophylaxis of the named disorder or condition, or amelioration or elimination of the disorder once it has been established.

"An effective amount" refers to an amount of a compound that confers a therapeutic effect on the treated subject. The therapeutic effect may be objective (i.e., measurable by some test or marker) or subjective (i.e., subject gives an indication of or feels an effect).

The term "prodrug forms" means a pharmacologically acceptable derivative, such as an ester or an amide, which derivative is biotransformed in the body to form the active drug. Reference is made to Goodman and Gilman's, *The Pharmacological basis of Therapeutics*, 8<sup>th</sup> ed., McGraw-Hill, Int. Ed. 1992, "Biotransformation of Drugs", p. 13-15; and "The Organic Chemistry of Drug Design and Drug Action" by Richard B. Silverman. Chapter 8, p 352. (Academic Press, Inc. 1992. ISBN 0-12-643730-0).

The following abbreviations have been used:

CV means Coefficient of Variation,  
 15 DMSO means dimethyl sulphoxide,  
 EDTA means ethylenediamine tetraacetic acid,  
 EGTA means ethylenebis(oxyethylenetrilo)tetraacetic acid,  
 HEPES means 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid,  
 HPLC means high performance liquid chromatography,  
 20 LSD means lysergic acid, diethylamide,  
 MeCN means acetonitrile,  
 SPA means Scintillation Proximity Assay,  
 t-BuOK means potassium tert-butoxide, and  
 THF means tetrahydrofuran.

25 All isomeric forms possible (pure enantiomers, diastereomers, tautomers, racemic mixtures and unequal mixtures of two enantiomers) for the compounds delineated are within the scope of the invention. Such compounds can also occur as cis- or trans-, *E*- or *Z*-double bond isomer forms. All isomeric forms are contemplated.

The compounds of the Formula (I) may be used as such or, where appropriate, as pharmacologically acceptable salts (acid or base addition salts) thereof. The pharmacologically acceptable addition salts mentioned above are meant to comprise the therapeutically active non-toxic acid and base addition salt forms that the compounds are able to form. Compounds that have basic properties can be converted to their pharmaceutically acceptable acid addition salts by treating the base form with an

appropriate acid. Exemplary acids include inorganic acids, such as hydrogen chloride, hydrogen bromide, hydrogen iodide, sulphuric acid, phosphoric acid; and organic acids such as formic acid, acetic acid, propanoic acid, hydroxyacetic acid, lactic acid, pyruvic acid, glycolic acid, maleic acid, malonic acid, oxalic acid, benzenesulphonic acid, 5 toluenesulphonic acid, methanesulphonic acid, trifluoroacetic acid, fumaric acid, succinic acid, malic acid, tartaric acid, citric acid, salicylic acid, p-aminosalicylic acid, pantoic acid, benzoic acid, ascorbic acid and the like. Exemplary base addition salt forms are the sodium, potassium, calcium salts, and salts with pharmaceutically acceptable amines such as, for example, ammonia, alkylamines, benzathine, and amino acids, such as, e.g. arginine 10 and lysine. The term addition salt as used herein also comprises solvates which the compounds and salts thereof are able to form, such as, for example, hydrates, alcoholates and the like.

For clinical use, the compounds of the invention are formulated into pharmaceutical formulations for oral, rectal, parenteral or other mode of administration. Pharmaceutical 15 formulations are usually prepared by mixing the active substance, or a pharmaceutically acceptable salt thereof, with conventional pharmaceutical excipients. Examples of excipients are water, gelatin, gum arabicum, lactose, microcrystalline cellulose, starch, sodium starch glycolate, calcium hydrogen phosphate, magnesium stearate, talcum, colloidal silicon dioxide, and the like. Such formulations may also contain other 20 pharmacologically active agents, and conventional additives, such as stabilizers, wetting agents, emulsifiers, flavouring agents, buffers, and the like. Usually, the amount of active compounds is between 0.1-95% by weight of the preparation, preferably between 0.2-20% by weight in preparations for parenteral use and more preferably between 1-50% by weight in preparations for oral administration.

25 The formulations can be further prepared by known methods such as granulation, compression, microencapsulation, spray coating, etc. The formulations may be prepared by conventional methods in the dosage form of tablets, capsules, granules, powders, syrups, suspensions, suppositories or injections. Liquid formulations may be prepared by dissolving or suspending the 30 active substance in water or other suitable vehicles. Tablets and granules may be coated in a conventional manner.

In a further aspect the invention relates to methods of making compounds of any of the formulae herein comprising reacting any one or more of the compounds of the



formulae delineated herein, including any processes delineated herein. The compounds of the formula (I) above may be prepared by, or in analogy with, conventional methods.

The processes described above may be carried out to give a compound of the invention in the form of a free base or as an acid addition salt. A pharmaceutically acceptable acid addition salt may be obtained by dissolving the free base in a suitable organic solvent and treating the solution with an acid, in accordance with conventional procedures for preparing acid addition salts from base compounds. Examples of addition salt forming acids are mentioned above.

The compounds of formula (I) may possess one or more chiral carbon atoms, and they may therefore be obtained in the form of optical isomers, e.g. as a pure enantiomer, or as a mixture of enantiomers (racemate) or as a mixture containing diastereomers. The separation of mixtures of optical isomers to obtain pure enantiomers is well known in the art and may, for example, be achieved by fractional crystallization of salts with optically active (chiral) acids or by chromatographic separation on chiral columns.

The chemicals used in the synthetic routes delineated herein may include, for example, solvents, reagents, catalysts, and protecting group and deprotecting group reagents. The methods described above may also additionally include steps, either before or after the steps described specifically herein, to add or remove suitable protecting groups in order to ultimately allow synthesis of the compounds. In addition, various synthetic steps may be performed in an alternate sequence or order to give the desired compounds. Synthetic chemistry transformations and protecting group methodologies (protection and deprotection) useful in synthesizing applicable compounds are known in the art and include, for example, those described in R. Larock, *Comprehensive Organic Transformations*, VCH Publishers (1989); T.W. Greene and P.G.M. Wuts, *Protective Groups in Organic Synthesis*, 3<sup>rd</sup> Ed., John Wiley and Sons (1999); L. Fieser and M. Fieser, *Fieser and Fieser's Reagents for Organic Synthesis*, John Wiley and Sons (1994); and L. Paquette, ed., *Encyclopedia of Reagents for Organic Synthesis*, John Wiley and Sons (1995) and subsequent editions thereof.


The necessary starting materials for preparing the compounds of formula (I) are either known or may be prepared in analogy with the preparation of known compounds. The dose level and frequency of dosage of the specific compound will vary depending on a variety of factors including the potency of the specific compound employed, the metabolic stability and length of action of that compound, the patient's age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the

severity of the condition to be treated, and the patient undergoing therapy. The daily dosage may, for example, range from about 0.001 mg to about 100 mg per kilo of body weight, administered singly or multiply in doses, e.g. from about 0.01 mg to about 25 mg each. Normally, such a dosage is given orally but parenteral administration may also be chosen.

The invention will now be further illustrated by the following non-limiting Examples.

The specific examples below are to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever. Without further elaboration, it is believed that one skilled in the art can, based on the description herein, utilize the present invention to its fullest extent. All publications cited herein are hereby incorporated by reference in their entirety.

TABLE 1




EXAMPLE	R <sup>1</sup>	R <sup>m</sup>
1		H
2		H
3		H
4		H

### Methods

<sup>1</sup>H nuclear magnetic resonance (NMR) and <sup>13</sup>C NMR were recorded on a Bruker Advance DPX 400 spectrometer at 400.1 and 100.6 MHz, respectively. All spectra were recorded using residual solvent or tetramethylsilane (TMS) as internal standard. Infra red (IR) spectra were recorded on a Perkin-Elmer Spectrum 1000 FT-IR spectrophotometer. Ionspray mass spectrometry (MS) spectra were obtained on a Perkin-Elmer API 150EX mass spectrometer. Accurate mass measurements were performed on a Micromass LCT dual probe. Preparative HPLC/MS was performed on a Waters/Micromass Platform ZQ system equipped with System A: ACE 5 C8 column (19x50mm), eluents: MilliQ water, MeCN and MilliQ/MeCN/0.1%TFA and system B: Xterra MS C18, 5µm column (19x50mm), eluents: MilliQ water, MeCN and NH<sub>4</sub>HCO<sub>3</sub> (100mM). Analytical HPLC were performed on Agilent 1100, column: ACE 3 C8 (system A) or column: YMC-Pack (system B), eluents: MilliQ/0.1%TFA and MeCN. Elemental analyses were performed on a Vario El instrument. Preparative flash chromatography was performed on Merck silica gel 60 (230-400 mesh).

## EXPERIMENTAL

### Synthesis of 4'-methyl-1',4',5',6'-tetrahydrospiro{piperidine-2,7'-pyrrolo[3,2-b]pyridine}




A mixture of 2-(2-ethylamino)pyrrole (Herz W. J. Am. Chem. Soc. 75, 483, 1953; Wasley J.W.F Eur. Pat.Appl. EP 338989 B, 1989) (9.6 g, 87 mmol) and 1-methylpiperazin-4-one (9.85 g, 87 mmol) in benzene (100 mL) was completely evaporated under vacuo. The dry residue was dissolved in methanol (50 mL). H<sub>2</sub>SO<sub>4</sub> (43 mL) in methanol (107 mL) was added to the methanol solution at 0 °C (ice). The cooling bath was removed and the mixture was stirred for 2 hours at room temperature. It resulted in the appearance of a white dense precipitate. Then the mixture was held overnight at -20 °C. The precipitate was filtered and dissolved in minimal amount of NaOH in water (30 %).

The organic material was extracted with ethyl acetate (100 mL x 4). The organic phases were combined and dried by  $K_2CO_3$  followed by filtration. The volatiles were eliminated under vacuo. The residue was triturated with cold ethyl acetate, filtered and washed by minimal amount of cold ethyl acetate to yield 7.6 g (44%) of product.

5

General method for the synthesis of 4'-methyl-1',4',5',6'-tetrahydrospiro{piperidine-2,7'-pyrrolo[3,2-b]pyridine}sulphonamides



10

A suspension of t-BuOK (160 mg, 1.4 mmol) in anhydrous THF (3 mL) was added to the mixture of 4'-methyl-1',4',5',6'-tetrahydrospiro{piperidine-2,7'-pyrrolo[3,2-b]pyridine} (205 mg, 1 mmol) in THF. The mixture was heated under stirring (complete dissolution was observed at 40 °C) followed by the addition of the corresponding sulfonyl chloride (1.2 mmol) in THF (3 mL). The mixture was rapidly heated to the boiling point and then cooled. The reaction mixture was poured into water, extracted with ethyl acetate, dried ( $K_2CO_3$ ) and evaporated. The product was isolated by column chromatography on silica gel ( $CHCl_3/MeOH$  :10/1). The yields of compounds synthesized by this method vary from 18 to 62%.

20

EXAMPLE 1

**4'-Methyl-1'-(2-naphthylsulphonyl)-1',4',5',6'-tetrahydrospiro{piperidine-2,7'-pyrrolo[3,2-b]pyridine} hydrochloride**

28.3 mg of material was isolated.  $^1H$  NMR (270 MHz, Methanol- $d_4$ )  $\delta$  2.25-2.50 (m, 4H), 2.96 (s, 3H), 3.19 (app. t., 2H), 3.52- 5.65 (m, 6H), 7.56 (d, 1H, J = 3.22 Hz), 7.72 (dq, 1H, J = 1.48 Hz), 7.83 (dd, 1H, J = 1.86 Hz, J = 8.98 Hz), 7.99-8.02 (m, 1H), 8.09 (d, 2H, J = 8.98 Hz), 8.11-8.15 (m, 1H), 8.67 (d, 1H, J = 1.73Hz). HPLC purity 96 %.

25

## EXAMPLE 2

**4'-Methyl-1'-(4-bromophenylsulphonyl)-1',4',5',6'-tetrahydrospiro{piperidine-2,7'-pyrrolo[3,2-b]pyridine} hydrochloride**

29.8 mg of material was isolated. <sup>1</sup>H NMR (270 MHz, DMSO-d<sub>6</sub>) δ 1.45-1.50 (bd, 2H) Hz), 1.71-1.83 (dt, 2H, J = 4.21 Hz), 2.15 (s, 3H), 2.36 (app.t, 2H), 2.45-2.55 (m, 2H), 2.59 (app.t, 2H), 2.88 (app.t, 2H), 2.86 (app.t, 2H), 6.32 (d, 1H, J = 3.46 Hz), 7.15 (d, 1H, J = 3.46 Hz), 7.43 (d, 1H, J = 3.96 Hz), 7.71 (d, 1H, J = 3.96 Hz). HPLC purity 95 %.

## EXAMPLE 3

10 **4'-Methyl-1'-(5-bromo-2-thienylsulphonyl)-1',4',5',6'-tetrahydrospiro{piperidine-2,7'-pyrrolo[3,2-b]pyridine} hydrochloride**

254 mg of material was isolated. <sup>1</sup>H NMR (270 MHz, DMSO-d<sub>6</sub>) δ 1.40-1.44 (bd, 2H, J = 12.62 Hz), 1.72 (dt, 2H, J = 4.21 Hz), 2.22 (s, 3H), 2.25 (bt, 2H, J = 10.89 Hz), 2.45-2.55 (m, 2H), 2.42-2.46 (m, 2H), 2.83 (app.t, 2H), 2.86 (app.t, 2H), 6.29 (d, 1H, J = 3.46 Hz), 7.15 (d, 1H, J = 3.22 Hz), 7.23 (dd, 1H, J = 4.95 Hz, J = 3.96 Hz), 7.83 (dd, 1H, J = 3.84 Hz, J = 1.36 Hz), 8.10 (dd, 1H, J = 4.95 Hz, J = 1.24 Hz). HPLC purity 95 %.

## EXAMPLE 4

20 **4'-Methyl-1'-(2-thienylsulphonyl)-1',4',5',6'-tetrahydrospiro{piperidine-2,7'-pyrrolo[3,2-b]pyridine} hydrochloride**

67.6 mg of material was isolated. <sup>1</sup>H NMR (270 MHz, Methanol-d<sub>4</sub>) δ 1.45 (d, 2H J = 13.11 Hz), 1.75 (dt, 2H, J = 3.96 Hz), 2.18 (s, 3H), 2.25-2.40 (m, 2H), 2.40-2.65 (m, 4H), 2.59 (app. t, 2H), 2.86 (app.t, 2H), 6.29 (d, 1H, J = 3.46 Hz), 7.15 (d, 1H, J = 3.22 Hz), 7.23 (dd, 1H, J = 4.95 Hz, J = 3.96 Hz), 7.83 (dd, 1H, J = 3.84 Hz, J = 1.36 Hz), 8.10 (dd, 1H, J = 4.95 Hz, J = 1.24 Hz). HPLC purity 95 %.

## BIOLOGICAL TESTS

30 The ability of a compound according to the invention to bind a 5-HT<sub>6</sub> receptor, and to be pharmaceutically useful, can be determined using *in vivo* and *in vitro* assays known in the art.

(a) 5-HT<sub>6</sub> binding Assay

Binding affinity experiment for the 5-HT<sub>6</sub> receptor are performed in HEK293 cells transfected with 5-HT<sub>6</sub> receptor using [<sup>3</sup>H]-LSD as labeled ligand according to the general method as described by Boess F.G et al. Neuropharmacology 36(4/5) 713-720, 1997.

5

## Materials

### Cell culture

The HEK-293 cell line transfected with the 5-HT<sub>6</sub> receptor was cultured in Dulbeccos Modified Eagles Medium containing 5 % dialyzed foetal bovine serum, (Gibco BRL 10106-169), 0.5 mM sodium pyruvate and 400 µg/ml Geneticin (G-418) (Gibco BRL10131-019). The cells were passaged 1:10, twice a week.

10

### Chemicals

The radioligand [<sup>3</sup>H] LSD 60-240 Ci/mmol, obtained from Amersham Pharmacia Biotech, (Buckinghamshire, England) was in ethanol and stored at -20°C. The compounds were dissolved in 100% DMSO and diluted with binding buffer.

15

### Disposable

Compounds were diluted in Costar 96 well V-bottom polypropylene plates (Corning Inc. Costar, NY, USA). Samples were incubated in Packard Optiplat (Packard Instruments B.V., Groningen, The Netherlands). The total amount of added radioligand was measured in Packard 24-well Barex plates (Packard Instruments B.V., Groningen, The Netherlands) in the presence of Microscint<sup>TM</sup> 20 scintillation fluid (Packard Bioscience, Meriden, CT, USA).

25

### Buffer

The binding buffer consisted of 20 mM HEPES, 150 mM NaCl, 10 mM MgCl<sub>2</sub>, and 1 mM, EDTA, pH 7.4.

30

### Methods

#### Membrane preparation

Cells were grown to approximately 90% confluence on 24.5 x 24.5 mm culture dishes. The medium was aspirated, and after rinsing with ice-cold PBS, the cells were scraped off

using 25 ml Tris buffer (50 mM Tris-HCl, 1 mM EDTA, 1 mM EGTA, pH 7.4) and a window scraper. The cells were then broken with a Polytron homogeniser, and remaining particulate matter was removed by low-speed centrifugation, 1000x g for 5 min. Finally, the membranes were collected by high-speed centrifugation (20 000x g), suspended in  
5 binding buffer, and frozen in aliquots at -70°C.

103000

### Radioligand binding

Frozen cell membranes were thawed, immediately rehomogenized with a Polytron homogenizer, and coupled to SPA wheat germ agglutinin beads (Amersham Life Sciences, Cardiff, England) for 30 min under continuous shaking of the tubes. After coupling, the  
 5 beads were centrifuged for 10 minutes at 1000 g, and subsequently suspended in 20 ml of binding buffer per 96-well plate. The binding reaction was then initiated by adding radioligand and test compounds to the bead-membrane suspension. Following incubation at room temperature, the assay plates were subjected to scintillation counting.

The original SPA method was followed except for that membranes were prepared from  
 10 HEK293 cells expressing the human 5-HT<sub>6</sub> receptor instead of from HeLa cells (Dinh DM, Zaworski PG, Gill GS, Schlachter SK, Lawson CF, Smith MW. Validation of human 5-HT<sub>6</sub> receptors expressed in HeLa cell membranes: saturation binding studies, pharmacological profiles of standard CNS agents and SPA development. (The Upjohn Company Technical Report 7295-95-064 1995;27 December). The specific binding of  
 15 [<sup>3</sup>H]-LSD was saturable, while the non-specific binding increased linearly with the concentration of added radioligand. [<sup>3</sup>H]-LSD bound with high affinity to 5-HT<sub>6</sub> receptors. The K<sub>d</sub> value was estimated to 2.6 ± 0.2 nM based on four separate experiments.

The total binding at 3 nM of [<sup>3</sup>H]-LSD, the radioligand concentration used in the  
 20 competition experiments, was typically 6000 dpm, and the specific binding more than 70%. 5-HT caused a concentration dependent inhibition of [<sup>3</sup>H]-LSD binding with an overall average K<sub>i</sub> value of 236 nM when tested against two different membrane preparations. The inter assay variability over three experiments showed a CV of 10% with an average K<sub>i</sub> values of 173 nM (SD 30) and a Hill coefficient of 0.94 (SD 0.09). The intra assay  
 25 variation was 3% (n=4). All unlabelled ligands displaced the specific binding of [<sup>3</sup>H]-LSD in a concentration-dependent manner, albeit at different potencies. The rank order of affinity for the 5-HT<sub>6</sub> receptor of reference compounds was methiothepin (K<sub>i</sub> 2 nM) > mianserin (190 nM) ≈ 5-HT (236 nM) > methysergide (482 nM) > mesulergine (1970 nM).

30

### Protein determination

Protein concentrations were determined with BioRad Protein Assay (Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing



the principle of protein-dye binding. Anal. Biochem. 1976;72:248-54). Bovine serum albumin was used as standard.

#### Scintillation counting

- 5 The radioactivity was determined in a Packard TopCount™ scintillation counter (Packard Instruments, Meriden, CT, USA) at a counting efficiency of approximately 20 %. The counting efficiency was determined in separate sets of experiments.

#### Saturation experiments

- 10 At least 6 concentrations in duplicates of radioligand (0.1-20 nM of [<sup>3</sup>H]-LSD) were used in saturation experiments. The specific binding was calculated as the difference between total binding and non-specific binding, which was determined as the binding of radioligand in the presence of 5 μM lisuride. B<sub>max</sub> and the dissociation constant, K<sub>d</sub>, were determined from the non-linear regression analysis using equation 1. L<sub>u</sub> is the unbound concentration  
15 of radioligand, and y is the amount bound.

$$y = \frac{B_{\max} \cdot Lu}{Lu + K_d} \quad (\text{equation 1})$$

#### Competition experiments

- Total- and non-specific binding of radioligand was defined in eight replicates of each.  
20 Samples containing test compound were run in duplicate at 11 concentrations. Incubations were carried out at room temperature for 3 hours. The IC<sub>50</sub> value, i.e. the concentration of test compound that inhibited 50% of the specific binding of radioligand, was determined with non linear regression analysis and the K<sub>i</sub> value was calculated using equation 2 [Cheng Y.C. Biochem. Pharmacol. 22, 3099-3108, 1973].

25 
$$K_i = \frac{IC_{50}}{1 + \frac{L}{K_d}} \quad (\text{equation 2})$$

L = concentration of radioligand

K<sub>d</sub> = Affinity of radioligand

#### *(b) 5-HT<sub>6</sub> Intrinsic Activity Assay*

Antagonists to the 5-HT<sub>6</sub> receptor were characterized by measuring inhibition of 5-HT induced increase in cAMP in HEK 293 cells expressing the human 5-HT<sub>6</sub> receptor (see Boess et al. (1997) *Neuropharmacology* 36: 713-720). Briefly, HEK293/5-HT<sub>6</sub> cells were seeded in polylysine coated 96-well plates at a density of 25,000 / well and grown in DMEM (Dulbecco's Modified Eagle Medium) (without phenol-red) containing 5% dialyzed Foetal Bovine Serum for 48 h at 37°C in a 5% CO<sub>2</sub> incubator. The medium was then aspirated and replaced by 0.1 ml assay medium (Hanks Balance Salt Solution containing 20 mM HEPES, 1.5 mM isobutylmethylxanthine and 1 mg/ml bovine serum albumin). After addition of test substances, 50 µl dissolved in assay medium, the cells were incubated for 10 min at 37°C in a 5% CO<sub>2</sub> incubator. The medium was again aspirated and the cAMP content was determined using a radioactive cAMP kit (Amersham Pharmacia Biotech, BIOTRAK RPA559). The potency of antagonists was quantified by determining the concentration that caused 50% inhibition of 5-HT (at [5-HT]= 8 times EC<sub>50</sub>) evoked increase in cAMP, using the formula  $IC_{50,corr} = IC_{50} / (1 + [5HT] / EC_{50})$ .

The compounds in accordance with the invention have a selective affinity to 5-HT<sub>6</sub> receptors with K<sub>i</sub> and IC<sub>50,corr</sub> values between 0.5 nM and 5 µM or display a % inhibition of [<sup>3</sup>H]-LSD ≥ 20 % at 50 nM and are antagonists, agonists or partial agonists at 5-HT<sub>6</sub>. The compounds show good selectivity over 5-HT<sub>1a</sub>, 5-HT<sub>1b</sub>, 5-HT<sub>2a</sub>, 5-HT<sub>2b</sub>, 5-HT<sub>2c</sub>.

TABLE 2

**Binding affinity (K<sub>i</sub>) at the 5-HT<sub>6</sub> receptor**

EXAMPLE	K <sub>i</sub> (nM)
1	140
3	160

(c) *In vivo* assay of reduction of food intake

For a review on serotonin and food intake, see Blundell, J.E. and Halford, J.C.G. (1998) *Serotonin and Appetite Regulation. Implications for the Pharmacological Treatment of Obesity*. *CNS Drugs* 9:473-495.

Obese (ob/ob) mouse is selected as the primary animal model for screening as this mutant mouse consumes high amounts of food resulting in a high signal to noise ratio. To further substantiate and compare efficacy data, the effect of the compounds on food consumption is also studied in wild type (C57BL/6J) mice. The amount of food consumed during 15 hours of infusion of compounds is recorded.

Male mice (obese C57BL/6JBom-Lep<sup>ob</sup> and lean wild-type C57BL/6JBom; Bomholtsgaard, Denmark) 8-9 weeks with an average body weight of 50 g (obese) and 25 g (lean) are used in all the studies. The animals are housed singly in cages at 23±1°C, 40-60 % humidity and have free access to water and standard laboratory chow. The 12/12-h light/dark cycle is set to lights off at 5 p.m. The animals are conditioned for at least one week before start of study.

The test compounds are dissolved in solvents suitable for each specific compound such as cyclodextrin, cyclodextrin/methane sulphonic acid, polyethylene glycol/methane sulphonic acid, saline. Fresh solutions are made for each study. Doses of 30, 50 and 100 mg kg<sup>-1</sup> day<sup>-1</sup> are used. The purity of the test compounds is of analytical grade.

The animals are weighed at the start of the study and randomized based on body weight. Alzet osmotic minipumps (Model 2001D; infusion rate 8 µl/h) are used and loaded essentially as recommended by the Alzet technical information manual (Alza Scientific Products, 1997; Theeuwes, F. and Yam, S.I. Ann. Biomed. Eng. 4(4). 343-353, 1976). Continuous subcutaneous infusion with 24 hours duration is used. The minipumps are either filled with different concentrations of test compounds dissolved in vehicle or with only vehicle solution and maintained in vehicle pre-warmed to 37°C (approx. 1h). The minipumps are implanted subcutaneously in the neck/back region under short acting anesthesia (metofane/enflurane). This surgical procedure lasts approximately 5 min.

The weight of the food pellets are measured at 5 p.m. and at 8 p.m. for two days before (baseline) and one day after the implantation of the osmotic minipumps. The weigh-in is performed with a computer assisted Mettler Toledo PR 5002 balance. Occasional spillage is corrected for. At the end of the study the animals are killed by neck dislocation and trunk blood sampled for later analysis of plasma drug concentrations.

The plasma sample proteins are precipitated with methanol, centrifuged and the supernatant is transferred to HPLC vials and injected into the liquid chromatography /mass spectrometric system. The mass spectrometer is set for electrospray positive ion mode and  
5 Multiple Reaction Monitoring. A linear regression analysis of the standards forced through the origin is used to calculate the concentrations of the unknown samples.


Food consumption for 15 hours is measured for the three consecutive days and the percentage of basal level values is derived for each animal from the day before and after  
10 treatment. The values are expressed as mean  $\pm$  SD and  $\pm$  SEM from eight animals per dose group. Statistical evaluation is performed by Kruskal-Wallis one-way ANOVA using the percent basal values. If statistical significance is reached at the level of  $p < 0.05$ , Mann-Whitney U-test for statistical comparison between control and treatment groups is  
15 performed.

The compounds according to the invention show an effect (i.e., reduction of food intake) in the range of 5-200 mg/kg/d.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

Claims


1. A compound of the Formula (I)




Formula (I)

wherein:

10 P is selected from a substituent of Formula (II) and Formula (III);



(II)



(III)

or P may also be selected from H or C<sub>1-6</sub>-alkyl provided that R<sup>m</sup> is selected from -NHSO<sub>2</sub>R<sup>11</sup>, -SO<sub>2</sub>NR<sup>8</sup>R<sup>11</sup> or -S(O)<sub>e</sub>R<sup>11</sup>, wherein e is 0, 1, 2 or 3;

v is 1 or 2;

----- represents a single bond or a double bond, with the proviso that both  
----- represent double bonds or that both ----- represent single bonds;

W<sub>1</sub>, W<sub>2</sub>, W<sub>3</sub>, Z and Y are each a carbon atom; or

20 one of W<sub>1</sub>, W<sub>2</sub>, W<sub>3</sub>, Z and Y is a nitrogen atom, while the remainder being carbon atoms, provided that both ----- in Formula (I) represent single bonds;

U is selected from CHR<sup>4</sup>, CR<sup>4</sup> and CR<sup>4</sup>R<sup>4'</sup>, provided that when the dotted line connecting W<sub>1</sub> and U is a double bond, then U is CR<sup>4</sup>; and further provided that when the dotted line connecting W<sub>1</sub> and U is a single bond, then U is selected from CHR<sup>4</sup> and CR<sup>4</sup>R<sup>4'</sup>;

$R^1$  is selected from:



- (a)  $C_{1-6}$ -alkyl,
- (b)  $C_{1-6}$ -alkoxy- $C_{1-6}$ -alkyl,
- (c)  $C_{2-6}$ -alkenyl,
- 5 (d) hydroxy- $C_{1-6}$ -alkyl,
- (e) halo- $C_{1-6}$ -alkyl,
- (f) aryl,
- (g) arylcarbonylmethyl,
- (h) aryl- $C_{2-6}$ -alkenyl,
- 10 (i) aryl- $C_{1-6}$ -alkyl,
- (j)  $C_{3-7}$ -cycloalkyl,
- (k) heteroaryl,
- (l) 4-piperidinyl,
- (m) N-substituted 4-piperidinyl, wherein the substituents are selected from  $C_{1-6}$ -alkyl, aryl-
- 15  $C_{1-6}$ -alkyl and heteroaryl- $C_{1-6}$ -alkyl,
- (n) heteroaryl- $C_{1-6}$ -alkyl,


wherein any heteroaryl or aryl residue, alone or as part of another group, is optionally substituted, independently, in one or more positions with substituents having the values as defined for  $R^m$  and  $R^{m'}$ ;

20


$R^m$  and  $R^{m'}$  are each independently selected from:

- (a) hydrogen,
- (b) halogen,
- (c)  $C_{1-6}$ -alkyl,
- 25 (d) hydroxy,
- (e)  $C_{1-6}$ -alkoxy,
- (f)  $C_{2-6}$ -alkenyl,
- (g) phenyl,
- (h) phenoxy,
- 30 (i) benzyloxy,
- (j) benzoyl,
- (k)  $-OCF_3$ ,
- (l)  $-CN$ ,

- (m) hydroxy-C<sub>1-6</sub>-alkyl,  
 (n) halo-C<sub>1-6</sub>-alkyl,  
 (o) -NR<sup>10</sup>R<sup>10</sup>,  
 (p) -NO<sub>2</sub>,  
 5 (q) -CONR<sup>10</sup>R<sup>10</sup>,  
 (r) -NHCO<sub>2</sub>R<sup>11</sup>,  
 (s) -NR<sup>8</sup>COR<sup>11</sup>,  
 (t) -SO<sub>2</sub>NR<sup>8</sup>R<sup>11</sup>,  
 (u) -C(=O)R<sup>11</sup>,  
 10 (v) C<sub>1-6</sub>-alkoxycarbonyl,  
 (w) -S(O)<sub>e</sub>R<sup>11</sup>, wherein e is 0, 1, 2 or 3,  
 (x) -SCF<sub>3</sub>,  
 (y) -CHF=CH<sub>2</sub>, or  
 (aa) -OCF<sub>2</sub>H;
- 15 and with the proviso that, when R<sup>m</sup> and R<sup>m'</sup> are different from hydrogen and ring A is aromatic, R<sup>m</sup> and R<sup>m'</sup> are attached to a carbon atom in Formula (I); and with the further proviso that when one of W<sub>1</sub>, W<sub>2</sub> and W<sub>3</sub> in Formula (I) is a nitrogen atom and both  represent single bonds the said nitrogen atom is attached to R<sup>m</sup>, wherein R<sup>m</sup> is selected from hydrogen or C<sub>1-4</sub>-alkyl and v is 1; and with the further proviso that when W<sub>1</sub>, W<sub>2</sub> and
- 20 W<sub>3</sub> in Formula (I) are each a carbon atom and both  represent single bonds, R<sup>m</sup> is selected from hydrogen or methyl; and with the further proviso that when R<sup>m</sup> or R<sup>m'</sup>, as substituents on ring A and B in Formula (I), are selected from phenyl, phenoxy, benzyloxy and benzoyl, the phenyl or aryl ring thereof may be optionally substituted by C<sub>1-4</sub>-alkyl, halogen, C<sub>1-4</sub>-alkoxy, C<sub>1-4</sub>-alkylthio, trifluoromethyl or cyano;
- 25 wherein R<sup>m</sup> and R<sup>4</sup> may be linked to each other to form a fused substituent of Formula (IV) provided that R<sup>m</sup> is attached to W<sub>1</sub>:



when U is CR<sup>4</sup> or CHR<sup>4</sup>, R<sup>4</sup> is a group selected from:





wherein  $R^4$  may also be a group selected from:  $O-(C=NH)NR^8R^8$  or  $-(CH_2)_t-O-NH(C=NH)-NR^8R^8$ ;

5 wherein:

$n = 0, 1, \text{ or } 2,$

$o = 1 \text{ or } 2,$

$p = 1, 2, 3, \text{ or } 4,$


$r = 2 \text{ or } 3,$

10

$s = 1, 2 \text{ or } 3,$

$t = 0, 1, 2, \text{ or } 3;$

when U is  $CHR^4$ ,  $R^4$  is additionally selected from the following groups:



15 wherein:

$n = 0, 1 \text{ or } 2,$

$o = 1 \text{ or } 2,$

$p = 1, 2, 3, \text{ or } 4,$

$r = 2 \text{ or } 3,$

20

$s = 1, 2 \text{ or } 3;$

wherein X is selected from O,  $NR^8$  and S;

wherein  $R^5$  is independently a group selected from:

25

(a) hydrogen,

- (b) C<sub>1-6</sub>-alkyl,  
 (c) 2-cyanoethyl,  
 (d) hydroxy-C<sub>2-6</sub>-alkyl,  
 (e) C<sub>3-6</sub>-alkenyl,  
 5 (f) C<sub>3-6</sub>-alkynyl,  
 (g) C<sub>3-7</sub>-cycloalkyl,  
 (h) C<sub>3-7</sub>-cycloalkyl-C<sub>1-4</sub>-alkyl,  
 (i) C<sub>1-6</sub>-alkoxy-C<sub>2-6</sub>-alkyl,  
 (j) aryl-C<sub>1-6</sub>-alkyl,  
 10 (k) heteroaryl-C<sub>1-6</sub>-alkyl,  
 (l) -C(=NH)-NR<sup>8</sup>R<sup>8</sup>,  
 (m) -C(=O)-NR<sup>8</sup>R<sup>8</sup>,  
 (n) -CH<sub>2</sub>-CO-NR<sup>8</sup>R<sup>8</sup>;
- 15 R<sup>6</sup> is selected from:  
 (a) hydrogen,  
 (b) C<sub>1-4</sub>-alkyl,  
 (c) hydroxy-C<sub>1-4</sub>-alkyl,  
 (d) C<sub>1-4</sub>-alkoxy-C<sub>1-4</sub>-alkyl,  
 20 (e) halo-C<sub>1-4</sub>-alkyl,  
 (f) -NR<sup>8</sup>R<sup>8</sup>,  
 (g) -CO-NR<sup>8</sup>R<sup>8</sup>;  
 (h) hydroxy, provided that the said hydroxy group is not attached to a carbon atom adjacent to a ring nitrogen atom;
- 25 R<sup>7</sup> is selected from:  
 (a) hydrogen,  
 (b) C<sub>1-4</sub>-alkyl,  
 (c) hydroxy-C<sub>1-4</sub>-alkyl, or  
 30 (d) C<sub>1-4</sub>-alkoxy-C<sub>1-4</sub>-alkyl,  
 (e) hydroxy, provided that the said hydroxy group is not attached to a carbon atom adjacent to a heterocyclic ring nitrogen atom and that the said hydroxy group is attached to a heterocyclic ring not substituted with oxo;

$R^8$  is each independently selected from:

- (a) hydrogen, or
- (b)  $C_{1-6}$ -alkyl,

5  $R^9$  is selected from:

- (a) hydrogen,
- (b)  $C_{1-4}$ -alkyl, wherein one or two groups may be present at any carbon atom, or when two groups are present at the same carbon atom they may together form a cyclopropane ring,
- 10 (c) hydroxy- $C_{1-4}$ -alkyl,
- (d)  $C_{1-4}$ -alkoxy- $C_{1-4}$ -alkyl,
- (e) halo- $C_{1-4}$ -alkyl,
- (f)  $-CO-NR^8R^8$ ;

15  $R^{10}$  is each independently selected from:

- (a) hydrogen,
- (b)  $C_{1-6}$ -alkyl
- (c)  $C_{3-7}$ -cycloalkyl, or

wherein the two  $R^{10}$  groups together with the nitrogen to which they are attached form a heterocyclic ring; and when the two  $R^{10}$  groups form a piperazine ring, the nitrogen of the piperazine ring that allows the substitution may be substituted with a group selected from  $R^5$ ;

20

$R^{11}$  is selected from:

- 25
- (a)  $C_{1-6}$ -alkyl,
  - (b) aryl, or
  - (c) heteroaryl;

$R^{12}$  is selected from:

- 30
- (a) hydrogen, or
  - (b) methyl;


when U is  $R^4R^{4'}$ ,  $R^4$  and  $R^{4'}$  are linked to each other to form a heterocyclic ring selected from pyrrolidine or piperidine, wherein the N atom may be substituted by a group selected

from  $R^5$ ; and pharmaceutically acceptable salts, hydrates, solvates, geometrical isomers, tautomers, optical isomers, and prodrug forms thereof.

2. The compound according to claim 1, wherein

5

P is selected from



(II)

each of  $W_1$ ,  $W_2$ ,  $W_3$ , Z and Y is a carbon atom provided that both  $\text{-----}$  in Formula (I) represent double bonds, or

10 one of  $W_1$ ,  $W_2$ ,  $W_3$ , Z and Y is a nitrogen atom, while the remainder being carbon atoms, provided that both  $\text{-----}$  in Formula (I) represent single bonds;

U is selected from  $\text{CHR}^4$ ,  $\text{CR}^4$  and  $\text{CR}^4\text{R}^{4'}$ , provided that when the dotted line connecting  $W_1$  and U is a double bond, then U is  $\text{CR}^4$ ; and further provided that when the dotted line  
15 connecting  $W_1$  and U is a single bond, then U is selected from  $\text{CHR}^4$  and  $\text{CR}^4\text{R}^{4'}$ ;

$R^1$  is selected from:

- (f) aryl,
- (h) aryl- $\text{C}_{2-6}$ -alkenyl,
- 20 (i) aryl- $\text{C}_{1-6}$ -alkyl,
- (j)  $\text{C}_{3-7}$ -cycloalkyl,
- (k) heteroaryl,
- (n) heteroaryl- $\text{C}_{1-6}$ -alkyl,

wherein any heteroaryl or aryl residue, alone or as part of another group, is optionally  
25 substituted, independently, in one or more positions with substituents having the values as defined for  $R^m$  and  $R^{m'}$ ;

$R^m$  and  $R^{m'}$  are each independently selected from:

- (a) hydrogen,
- 30 (b) halogen,


- (c) C<sub>1-6</sub>-alkyl,
- (d) hydroxy,
- (e) C<sub>1-6</sub>-alkoxy,
- (f) C<sub>2-6</sub>-alkenyl,
- 5 (k) -OCF<sub>3</sub>,
- (m) hydroxy-C<sub>1-6</sub>-alkyl,
- (n) halo-C<sub>1-6</sub>-alkyl,
- (o) -NR<sup>10</sup>R<sup>10</sup>,
- (q) -CONR<sup>10</sup>R<sup>10</sup>,
- 10 (r) -NHSO<sub>2</sub>R<sup>11</sup>,
- (s) -NR<sup>8</sup>COR<sup>11</sup>,
- (t) -SO<sub>2</sub>NR<sup>8</sup>R<sup>11</sup>,
- (u) -C(=O)R<sup>11</sup>,
- (w) -S(O)<sub>e</sub>R<sup>11</sup>, wherein e is 0, 1, 2 or 3,
- 15 (x) -SCF<sub>3</sub>,
- (y) -CHF=CH<sub>2</sub>, or
- (aa) -OCF<sub>2</sub>H;

and with the proviso that, when R<sup>m</sup> and R<sup>m'</sup> are different from hydrogen and ring A is aromatic, R<sup>m</sup> and R<sup>m'</sup> are attached to a carbon atom in Formula (I); and with the further

20 proviso that when one of W<sub>1</sub>, W<sub>2</sub> and W<sub>3</sub> in Formula (I) is a nitrogen atom and both        represent single bonds the said nitrogen atom is attached to R<sup>m</sup>, wherein R<sup>m</sup> is selected from hydrogen or C<sub>1-4</sub>-alkyl and v is 1; and with the further proviso that when W<sub>1</sub>, W<sub>2</sub> and W<sub>3</sub> in Formula (I) are each a carbon atom and both        represent single bonds, R<sup>m</sup> is selected from hydrogen or methyl; and with the further proviso that when R<sup>m</sup> or R<sup>m'</sup>, as

25 substituents on ring A and B in Formula (I), are selected from phenyl, phenoxy, benzyloxy and benzoyl, the phenyl or aryl ring thereof may be optionally substituted by C<sub>1-4</sub>-alkyl, halogen, C<sub>1-4</sub>-alkoxy, C<sub>1-4</sub>-alkylthio, trifluoromethyl or cyano;

when U is CR<sup>4</sup> or CHR<sup>4</sup>, R<sup>4</sup> is a group selected from:




$n = 0, 1, \text{ or } 2,$

$o = 1 \text{ or } 2,$

$p = 1, 2, 3, \text{ or } 4,$

5

when U is CHR<sup>4</sup>, R<sup>4</sup> is additionally selected from the following groups:



wherein :

$n = 0, 1 \text{ or } 2,$

$o = 1 \text{ or } 2,$

5

$p = 1, 2, \text{ or } 3;$

$r = 2 \text{ or } 3,$

wherein X is selected from O,  $\text{NR}^8$  and S;

wherein  $\text{R}^5$  is independently a group selected from:

10

(a) hydrogen,

(b)  $\text{C}_{1-6}$ -alkyl,

(c) 2-cyanoethyl,

(d) hydroxy- $\text{C}_{2-4}$ -alkyl,

(e)  $\text{C}_{3-6}$ -alkenyl,

15

(h)  $\text{C}_{3-7}$ -cycloalkyl- $\text{C}_{1-4}$ -alkyl,

(i)  $\text{C}_{1-4}$ -alkoxy- $\text{C}_{2-4}$ -alkyl,

$\text{R}^6$  is selected from:

(a) hydrogen,

20

(b)  $\text{C}_{1-4}$ -alkyl,

(f)  $-\text{NR}^8\text{R}^8,$

$\text{R}^7$  is selected from:

(a) hydrogen,

25

(b)  $\text{C}_{1-4}$ -alkyl,

(c) hydroxy-C<sub>1-2</sub>-alkyl, or

(d) C<sub>1-2</sub>-alkoxy-C<sub>1-2</sub>-alkyl,

(e) hydroxy, provided that the said hydroxy group is not attached to a carbon atom adjacent to a heterocyclic ring nitrogen atom and that the said hydroxy group is attached to a heterocyclic ring not substituted with oxo;

R<sup>8</sup> is each independently selected from:

(a) hydrogen, or

(b) C<sub>1-6</sub>-alkyl,

R<sup>9</sup> is selected from:

(a) hydrogen,

(b) C<sub>1-4</sub>-alkyl, wherein one or two groups may be present at any carbon atom, or when two groups are present at the same carbon atom they may together form a cyclopropane ring,

(c) hydroxy-C<sub>1-2</sub>-alkyl,

(d) C<sub>1-2</sub>-alkoxy-C<sub>1-2</sub>-alkyl,

(e) halo-C<sub>1-2</sub>-alkyl,

R<sup>10</sup> is each independently selected from:

(a) hydrogen,

(b) C<sub>1-6</sub>-alkyl, or

wherein the two R<sup>10</sup> groups together with the nitrogen to which they are attached form a heterocyclic ring; and when the two R<sup>10</sup> groups form a piperazine ring, the nitrogen of the piperazine ring that allows the substitution may be substituted with a group selected from R<sup>5</sup>;

R<sup>11</sup> is selected from:

(a) C<sub>1-6</sub>-alkyl


provided that P is a group selected from Formula (II) or Formula (III);

R<sup>12</sup> is selected from:


(a) hydrogen, or

(b) methyl;





(II)



(III)

when U is  $R^4R^{4'}$ ,  $R^4R^{4'}$  is linked to each other to form a heterocyclic ring selected from pyrrolidine or piperidine, wherein the N atom may be substituted by a group  $R^5$  selected from:

- (a) hydrogen,
- (b)  $C_{1-6}$ -alkyl,
- (c) 2-cyanoethyl,
- (d) hydroxy- $C_{2-4}$ -alkyl,
- (i)  $C_{1-4}$ -alkoxy- $C_{2-4}$ -alkyl.

3. The compound according to any one of claims 1 or 2, wherein

ring A is saturated;

15  $W_1$  is a nitrogen atom;

Z and Y are both carbon atoms;

$R^1$  is a group aryl or heteroaryl optionally substituted by halogen;

$R^m$  and  $R^{m'}$  are both H;

wherein  $R^4$  and  $R^{4'}$  are linked to each other to form piperidine, wherein the N atom is substituted by methyl.

4. The compound according to any one of claims 1 to 3, wherein

Ar is selected from phenyl, naphthyl, and thienyl, which group Ar is optionally substituted by halogen.

5. The compound according to any one of claims 1 to 4, which is selected from 4'-Methyl-1'-(2-naphthylsulphonyl)-1',4',5',6'-tetrahydrospiro {piperidine-2,7'-pyrrolo[3,2-b]pyridine} hydrochloride,

4'-Methyl-1'-(4-bromophenylsulphonyl)-1',4',5',6'-tetrahydrospiro {piperidine-2,7'-pyrrolo[3,2-b]pyridine} hydrochloride,

4'-Methyl-1'-(5-bromo-2-thienylsulphonyl)-1',4',5',6'-tetrahydrospiro {piperidine-2,7'-pyrrolo[3,2-b]pyridine} hydrochloride, and

4'-Methyl-1'-(2-thienylsulphonyl)-1',4',5',6'-tetrahydrospiro {piperidine-2,7'-pyrrolo[3,2-b]pyridine} hydrochloride.

5

6. A process for the preparation of a compound according to any one of claims 1 to 5, comprising the following steps:

a) reaction of 2-(2-ethylamino)pyrrole and 1-methylpiperazine-4-one to give 4'-methyl-1',4',5',6'-tetrahydrospiro {piperidine-2,7'-pyrrolo[3,2-b]pyridine}; and

10 b) reaction of the product from step a) with an arylsulphonyl chloride in the presence of a base.

7. A compound according to any one of claims 1 to 5 for use in therapy.

15 8. A compound according to any one of claims 1 to 5 for use in the treatment or prophylaxis of a 5-HT<sub>6</sub> receptor-related disorder, to achieve reduction of body weight and of body weight gain.

20 9. A compound according to claim 8, wherein the disorder is selected from obesity; type II diabetes; disorders of the central nervous system such as anxiety, depression, panic attacks, memory disorders, cognitive disorders, epilepsy, sleep disorders, migraine, anorexia, bulimia, binge eating disorders, obsessive compulsive disorders, psychoses, Alzheimer's disease, Parkinson's disease, Huntington's chorea, schizophrenia, attention deficit hyperactive disorder, withdrawal from drug abuse, neurodegenerative diseases  
25 characterized by impaired neuronal growth, and pain.

30 10. A pharmaceutical formulation comprising a compound according to any one of claims 1 to 5 as active ingredient, in combination with a pharmaceutically acceptable diluent or carrier.

11. The pharmaceutical formulation according to claim 10 for use in the prophylaxis or treatment of a 5-HT<sub>6</sub> receptor-related disorder, to achieve reduction of body weight and of body weight gain.

12. The pharmaceutical formulation according to any one of claim 10 or 11, wherein the disorder is selected from obesity; type II diabetes; disorders of the central nervous system such as anxiety, depression, panic attacks, memory disorders, cognitive disorders, epilepsy, sleep disorders, migraine, anorexia, bulimia, binge eating disorders, obsessive compulsive disorders, psychoses, Alzheimer's disease, Parkinson's disease, Huntington's chorea, schizophrenia, attention deficit hyperactive disorder, withdrawal from drug abuse, neurodegenerative diseases characterized by impaired neuronal growth, and pain.
13. A method for the prophylaxis or treatment of a 5-HT<sub>6</sub> receptor-related disorder, to achieve reduction of body weight and of body weight gain, which comprises administering to a subject in need of such treatment an effective amount of a compound according to any one of claims 1 to 5.
14. The method according to claim 13, wherein the disorder is selected from obesity; type II diabetes; disorders of the central nervous system such as anxiety, depression, panic attacks, memory disorders, cognitive disorders, epilepsy, sleep disorders, migraine, anorexia, bulimia, binge eating disorders, obsessive compulsive disorders, psychoses, Alzheimer's disease, Parkinson's disease, Huntington's chorea, schizophrenia, attention deficit hyperactive disorder, withdrawal from drug abuse, neurodegenerative diseases characterized by impaired neuronal growth, and pain.
15. A method for modulating 5-HT<sub>6</sub> receptor activity, which comprises administering to a subject in need of such treatment an effective amount of a compound according to any one of claims 1 to 5.
16. Use of a compound according to any one of claims 1 to 5 for the manufacture of a medicament for use in the prophylaxis or treatment of a 5-HT<sub>6</sub> receptor-related disorder, to achieve reduction of body weight and of body weight gain.
17. The use according to claim 16, wherein the disorder is selected from obesity; type II diabetes; disorders of the central nervous system such as anxiety, depression, panic attacks, memory disorders, cognitive disorders, epilepsy, sleep disorders, migraine, anorexia,

5


18. A cosmetic composition comprising a compound according to any one of claims 1 to 5 as active ingredient, in combination with a cosmetically acceptable diluent or carrier.

19. The cosmetic composition according to claim 18 for use in the prophylaxis or treatment  
10 of a 5-HT<sub>6</sub> receptor-related disorder, to achieve reduction of body weight and of body  
weight gain.

10

## **Abstract**

**The present invention relates to compounds of formula (I):**



### Formula (I)

wherein

U, P, W<sub>1</sub>, W<sub>2</sub>, W<sub>3</sub>, v, Y, Z, R<sup>m</sup>, and R<sup>m'</sup> are as described herein, to pharmaceutical compositions comprising the compounds, to processes for their preparation, as well as to the use of the compounds for the preparation of a medicament against 5-HT<sub>6</sub> receptor-related disorders.